

FERTIACTYL[®] ORGANIC

NON PLANT FOOD INGREDIENTS:
Humic acid.....4%
Fulvic acid.....1%
Seaweed extract.....20%

DIRECTIONS FOR USE:

- FERTIACTYL[®] Organic may be used in application to the soil, near the root system. Fertilization system or other ways of spreading the product can be used, like overhead irrigation.
- Do not spray during periods of frost. In hot conditions it is better to spray at the beginning or at the end of the day. Fill tank half full with clean water then add FERTIACTYL[®] Organic and continue to agitate while mixing. Keep agitating until the end of the application. Dilute into at least 20 Gal/Acre.
- FERTIACTYL[®] Organic is intended as a supplement to a regular fertilization program and may not by itself provide all the nutrients normally required by crops or other intended plants.

STORAGE:

- Do not store under 32°F (0°C).

SHAKE WELL BEFORE USE

NET VOLUME: 1 Gal (3.79 L) - NET WEIGHT: 8.67 lb (3.93 kg)

Timac Agro USA, Inc. warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for purposes stated on the label when used in accordance with directions under normal use conditions. Crop Injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Timac Agro USA, Inc. In no case shall Timac Agro USA, Inc. be liable for consequential, special or indirect damages resulting from the use, handling, or shipping of this product. All such risk shall be assumed by the Buyer. Timac Agro USA, Inc. makes no warranties of merchantability or fitness for a particular purpose nor any other express or implied warranty except as stated above.



MANUFACTURED BY
TIMAC AGRO USA, INC
P.O. Box 888
Route 724 & I-176
Reading, PA 19607-0888 -USA

Guaranteed Analysis:

Nitrogen (N).....0%
Soluble Phosphorus (P205)....0%
Soluble Potash (K2O).....1%

Derived from:

Seaweed extract



KEEP OUT OF
REACH OF
CHILDREN

F1550

APPLICATION TIMING CHART

Crop	Nº of App.	Application Stage	App Rate (pts/Acre)	Advice
Corn, Soybean & Alfalfa	1-4	At planting	2-4	Broadcast app.
			15-4	In row app.
	Early growth	2-4	Broadcast app.	
		15-4	Banded app.	
Orchard/ Vineyard	1-4	At planting	0.25-1% transplant solution	4 pts/Acre gives best results.
		Bloom to Pre-Harvest	2-4	
General Vegetables	1-4	At planting	0.25-1% transplant solution	Must have mechanical or by pass tank agitation in transplant water tank
		At transplant to 7 days after planting	2-4	
		With 2nd foliar spray (7-14 days after 1st application)		

FERTIACTYL[®] ORGANIC

NON PLANT FOOD INGREDIENTS:
Humic acid.....4%
Fulvic acid.....1%
Seaweed extract.....20%

DIRECTIONS FOR USE:

- FERTIACTYL[®] Organic may be used in application to the soil, near the root system. Fertilization system or other ways of spreading the product can be used, like overhead irrigation.
- Do not spray during periods of frost. In hot conditions it is better to spray at the beginning or at the end of the day. Fill tank half full with clean water then add FERTIACTYL[®] Organic and continue to agitate while mixing. Keep agitating until the end of the application. Dilute into at least 20 Gal/Acre.
- FERTIACTYL[®] Organic is intended as a supplement to a regular fertilization program and may not by itself provide all the nutrients normally required by crops or other intended plants.

STORAGE:

- Do not store under 32°F (0°C).

SHAKE WELL BEFORE USE

NET VOLUME: 2.64 Gal (10L) - NET WEIGHT: 22.89 lb (10.38 kg)

Timac Agro USA, Inc. warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for purposes stated on the label when used in accordance with directions under normal use conditions. Crop Injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Timac Agro USA, Inc. In no case shall Timac Agro USA, Inc. be liable for consequential, special or indirect damages resulting from the use, handling, or shipping of this product. All such risk shall be assumed by the Buyer. Timac Agro USA, Inc. makes no warranties of merchantability or fitness for a particular purpose nor any other express or implied warranty except as stated above.



MANUFACTURED BY
TIMAC AGRO USA, INC
P.O. Box 888
Route 724 & I-176
Reading, PA 19607-0888 -USA

Guaranteed Analysis:

Nitrogen (N).....0%
Soluble Phosphorus (P205)....0%
Soluble Potash (K2O).....1%

Derived from:

Seaweed extract



KEEP OUT OF
REACH OF
CHILDREN

F1550

APPLICATION TIMING CHART

Crop	Nº of App.	Application Stage	App Rate (pts/Acre)	Advice	
Corn, Soybean & Alfalfa	1-4	At planting	2-4	Broadcast app.	
			15-4	In row app.	
	Early growth	2-4	Broadcast app.		
		15-4	Banded app.		
Orchard/ Vineyard	1-4	At planting	0.25-1% transplant solution	4 pts/Acre gives best results.	
		Bloom to Pre-Harvest	2-4		
General Vegetables	1-4	At planting	0.25-1% transplant solution	Must have mechanical or by pass tank agitation in transplant water tank	
		At transplant to 7 days after planting	2-4		Injected in clean water (drip irrigation). If not available, use it to water plants (same rate) wetting Leaves until the solution reaches the soil.
		With 2nd foliar spray (7-14 days after 1st application)			

FERTIACTYL[®] ORGANIC

NON PLANT FOOD INGREDIENTS:
Humic acid.....4%
Fulvic acid.....1%
Seaweed extract.....20%

DIRECTIONS FOR USE:

- FERTIACTYL[®] Organic may be used in application to the soil, near the root system. Fertilization system or other ways of spreading the product can be used, like overhead irrigation.
- Do not spray during periods of frost. In hot conditions it is better to spray at the beginning or at the end of the day. Fill tank half full with clean water then add FERTIACTYL[®] Organic and continue to agitate while mixing. Keep agitating until the end of the application. Dilute into at least 20 Gal/Acre.
- FERTIACTYL[®] Organic is intended as a supplement to a regular fertilization program and may not by itself provide all the nutrients normally required by crops or other intended plants.

STORAGE:

- Do not store under 32°F (0°C).

SHAKE WELL BEFORE USE

NET VOLUME: 264Gal (999.35L) - NET WEIGHT: 2289lb (1038.27kg)

Timac Agro USA, Inc. warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for purposes stated on the label when used in accordance with directions under normal use conditions. Crop Injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Timac Agro USA, Inc. In no case shall Timac Agro USA, Inc. be liable for consequential, special or indirect damages resulting from the use, handling, or shipping of this product. All such risk shall be assumed by the Buyer. Timac Agro USA, Inc. makes no warranties of merchantability or fitness for a particular purpose nor any other express or implied warranty except as stated above.



MANUFACTURED BY
TIMAC AGRO USA, INC
P.O. Box 888
Route 724 & I-176
Reading, PA 19607-0888 -USA

Guaranteed Analysis:

Nitrogen (N).....0%
Soluble Phosphorus (P205)....0%
Soluble Potash (K2O).....1%

Derived from:

Seaweed extract



KEEP OUT OF
REACH OF
CHILDREN

F1550

APPLICATION TIMING CHART

Crop	Nº of App.	Application Stage	App Rate (pts/Acre)	Advice	
Corn, Soybean & Alfalfa	1-4	At planting	2-4	Broadcast app.	
			15-4	In row app.	
	Early growth	2-4	Broadcast app.		
		15-4	Banded app.		
Orchard/ Vineyard	1-4	At planting	0.25-1% transplant solution	4 pts/Acre gives best results.	
		Bloom to Pre-Harvest	2-4		
General Vegetables	1-4	At planting	0.25-1% transplant solution	Must have mechanical or by pass tank agitation in transplant water tank	
		At transplant to 7 days after planting	2-4		Injected in clean water (drip irrigation). If not available, use it to water plants (same rate) wetting Leaves until the solution reaches the soil.
		With 2nd foliar spray (7-14 days after 1st application)			